KNEE



Unicompartmental knee arthroplasty versus total knee arthroplasty: Which type of artificial joint do patients forget?

Hendrik A. Zuiderbaan¹ · Jelle P. van der List¹ · Saker Khamaisy¹ · Danyal H. Nawabi¹ · Ran Thein¹ · C. Ishmael¹ · Sophia Paul¹ · Andrew D. Pearle¹

Received: 10 August 2015 / Accepted: 5 November 2015 © European Society of Sports Traumatology, Knee Surgery, Arthroscopy (ESSKA) 2015

Abstract

Purpose During recent years, there has been an intensive growth of interest in the patient's perception of functional outcome. The Forgotten Joint Score (FJS) is a recently introduced score that measures joint awareness of patients who have undergone knee arthroplasty and is less limited by ceiling effects. The aim of this study was to compare the FJS between patients who undergo medial unicompartmental knee arthroplasty (UKA) and patients who undergo total knee arthroplasty (TKA) 1 and 2 years post-operatively.

Methods This prospective study compares the FJS at a minimum of one (average 1.5 years, range 1.0–1.9) and a minimum of 2 years (average 2.5 years, range 2.0–3.6)

\bowtie	Hendrik A. Zuiderbaan
	hazuiderbaanmd@gmail.com

Jelle P. van der List vanderlistj@hss.edu

Saker Khamaisy Saker.khamaisy@gmail.com

Danyal H. Nawabi nawabid@hss.edu

Ran Thein ranthein@gmail.com

C. Ishmael cishmael@gmail.com

Sophia Paul pauls@hss.edu

Andrew D. Pearle pearlea@hss.edu

¹ Department of Orthopaedic Surgery, Computer Assisted Surgery Center, Hospital for Special Surgery, Weill Medical College of Cornell University, 535 E. 70th Street, New York, NY 10021, USA post-operatively between patients who underwent medial UKA and TKA.

Results One-hundred and thirty patients were included. Sixty-five patients underwent medial UKA and 65 patients underwent TKA. At both follow-up points, the FJS was significantly higher in the UKA group (FJS 1 year 73.9 \pm 22.8, FJS 2 year 74.3 \pm 24.8) in contrast to the TKA group (FJS 1 year 59.3 \pm 29.5 (p = 0.002), FJS 2 year 59.8 \pm 31.5, (p = 0.004)). No significant improvement in the FJS was observed between 1- and 2-year follow-up of the two cohorts.

Conclusion Patients who undergo UKA are more likely to forget their artificial joint in daily life and consequently may be more satisfied.

Level of evidence II.

Keywords Unicompartmental knee arthroplasty · Total knee arthroplasty · Forgotten joint score · Subjective outcome

Introduction

Total knee replacement surgery is the accepted treatment for end-stage arthritis of the knee. In 2008, more than 600,000 cases were performed in the USA [28]. Recently, the utilization of unicompartmental knee arthroplasty (UKA) has dramatically increased for end-stage arthritis of the knee localized to a single compartment [12, 18, 20]. Less perioperative blood loss, better range of motion, better quadriceps function and a subsequently quicker recovery with a more normal gait [5] following UKA may all have contributed to the increasing utilization of the implant.

As a result of the reported advantages of UKA, there has been increasing interest in comparing the outcomes of

UKA and total knee arthroplasty (TKA), with a particular focus on patients' perception of functional outcome. The outcomes of joint replacement have historically been evaluated based on implant survivorship, physician-assessed clinical outcome measures, complication rates and radiological parameters. Although these outcomes are critically important to report, they do not provide any information pertaining to patient perception of outcome. Therefore, patient-reported outcome (PRO) scores were developed and validated for clinical use. These scores are, however, limited firstly by ceiling effects [15], particularly in young and active patients, and secondly by the heterogeneity of scores in current use, making it difficult to compare outcomes with previously published data [2, 13, 14, 19, 24].

The Forgotten Joint Score (FJS) [3] is a recently validated PRO score which is not limited by a ceiling effect [27]. The rationale thought behind the FJS was to develop an instrument that reflects the ability of a patient to perform activities of daily living (ADL) without any form of interference from their artificial joint replacement. Since the FJS is a relatively new score, prospective data of patients undergoing knee arthroplasty using the FJS are very scarce. Therefore, the aim of this study was to compare the FJS between patients undergoing either medial UKA or TKA in order to evaluate the artificial joint awareness in both groups at 1 and 2 years following surgery.

Materials and methods

This study was based on a prospective cohort of patients assembled for the senior author's surgical arthritis registry. Patients were eligible for this analysis if they were adult participants in the registry and underwent medial UKA or TKA between 2011 and 2013. Patient demographics and clinical data were collected including age, body mass index (BMI) and gender. The surgical indications of patients who underwent medial UKA consisted of: (I) isolated medial compartment OA, (II) an intact anterior cruciate ligament based on clinical and intraoperative assessments, (III) flexion contracture $<10^{\circ}$ and (IV) $>90^{\circ}$ of tibiofemoral flexion. The indications for TKA were (I) symptomatic OA changes of at least two compartments of the knee and (II) symptomatic OA of medial or lateral compartment of the knee in patients who did not wish to undergo UKA or in patients with proven anterior cruciate ligament deficiency. The presence of anterior knee pain and pre-operative Kellgren and Lawrence grade III-IV of the lateral or patellofemoral compartments was considered as surgical contraindications for medial UKA, and those patients therefore underwent TKA. Patients with a history of complex knee surgery, trauma, inflammatory arthropathy, BMI >40 kg/m² and simultaneous bilateral TKA or UKA were excluded.

All surgeries were performed by the senior author who has extensive experience in computer-navigated robotic surgery. For the UKA resurfacing procedure of the femur and tibia, a robotic arm-assisted technique was used, which has previously been described [21, 23] (MAKO Surgical Corp, Ft. Lauderdale, FL, USA). The goal was an undercorrection of the varus deformity in order to avoid degenerative progression of the lateral compartment. All patients who underwent medial UKA received the RESTORIS[®] MCK Medial Onlay implant (MAKO Surgical Corporation, Ft. Lauderdale, FL, USA).

All patients who underwent TKA received the Vanguard[®] Complete Total Knee (Biomet, Warsaw, USA) utilizing patient-specific cutting jigs based on pre-operative computed tomography. All procedures were performed under tourniquet control, with patellar resurfacing, and cementation of all implants.

Outcome measurements

All patients were asked to complete the FJS at a minimum of 1- and 2-year follow-up. The date of the 2-year data collection had to be at least 12 months later than the date of the 1-year data collection. Since the FJS is a questionnaire following arthroplasty, it is not possible to collect pre-operative scores (Appendix). The FJS consists of 12 questions. It has been devised to evaluate the ability of a patient to forget their artificial joint in daily life. The score is reported on a scale from 0 to 100. A higher score is representative of a more favourable outcome. A detailed description on how the score should be calculated is given in the "Appendix".

This study was approved by the institutional review board at our hospital (Hospital for Special Surgery, New York, NY, USA. IRB number: 2013-056-CR2).

Statistical analysis

All analyses were conducted using SAS for Windows 9.3 (SAS Institute Inc., Cary, NC). A priori power analysis was conducted using a two-sample *t* test. Sixty-four patients in each group were needed to reach 80 % power for detecting a 12-point (standard deviation 24) difference on the FJS scale with a two-sided significance level set at 0.05. Two-sample *t* tests were used to compare the FJS results between the medial UKA and TKA groups and between male and female subjects. A multiple linear regression analysis was performed to assess the relationship between surgery type and FJS score, controlling for age, sex, BMI and duration of follow-up. All tests were two-sided. A *p* value <0.05 was considered as statistically significant.

 Table 1
 Baseline

 characteristics
 Image: Characteristic state

	Medial UKA			ТКА			
	N	Mean	SD	N	Mean	SD	p value
Female sex. N (%)	28 (43.1 %)			37 (56.9 %)			n.s.
Age (years)		66.6	10.5		67.9	8.4	n.s.
BMI (kg/m ²)		28.6	3.7		30.3	4.7	0.02
1 year F/U (range)		1.5 (1.0–1.9)	0.4		1.4 (1.0–1.9)	0.3	n.s.
2 year F/U (range)		2.4 (2.0-3.4)	0.7		2.6 (2.0-3.6)	0.6	n.s.

Results

One-hundred and thirty patients were able to complete the outcome questionnaires at a minimum of 1-year followup. Both groups consisted of 65 patients (Table 1); 2 years following surgery, three patients were lost to follow-up (1 medial UKA, 2 TKA) because they moved away from the area. There were no significant differences in age, gender distribution and average follow-up at 1- and 2-year follow-up between the two cohorts. The average BMI in the medial UKA group (28.6 \pm 3.7 kg/m²) was significantly lower than in the TKA group (30.3 \pm 4.7 kg/m²) (p = 0.02) (Table 1). During the follow-up none of the included patients were re-operated or underwent revision surgery.

Outcome measurements

One year following surgery, the mean FJS in the medial UKA group (73.9 \pm 22.8) was significantly higher than the TKA group (59.3 \pm 29.5, p = 0.002); 2 years following surgery, the FJS remained significantly higher in the medial UKA group (74.3 \pm 24.8) in comparison with the TKA group (59.8 \pm 31.5, p = 0.004) (Fig. 1). No significant improvement in the two groups was observed over time when comparing the 1- and 2-year FJS data (Table 2).

No significant differences were found in the FJS between men and women in both the medial UKA and TKA groups at 1 and 2 years following surgery (Table 3). Multivariate regression analysis showed the FJS in the medial UKA group 1 year following surgery to be significantly higher than the TKA group (72.2 vs. 61.1, p = 0.02), after controlling for BMI, age, sex and follow-up. This significant difference of the FJS remained at 2-year follow-up in favour of the medial UKA group (72.4 vs. 61.2, p = 0.01).

Discussion

The most important finding of the present study is that patients who have undergone medial UKA are less aware of their artificial joint than patients who have undergone TKA. The FJS has the ability to distinguish between good and excellent outcomes and is therefore not limited



Fig. 1 Forgotten Joint Score 1 and 2 years following surgery. Note that the medial unicompartmental knee arthroplasty (UKA) group showed significant higher scores at 1- and 2-year follow-up (FJS 1 year 73.9 \pm 22.8, FJS 2 year 74.3 \pm 24.8) in contrast to the total knee arthroplasty (TKA) group (FJS 1 year 59.3 \pm 29.5, p = 0.002), FJS 2 year 59.8 \pm 31.5, p = 0.004)

by ceiling effects. There is no consensus in the literature regarding the efficacy of medial UKA compared to TKA with respect to patient satisfaction. The purpose of this study was to compare outcomes of medial UKA and TKA using the FJS at a minimum of 1- and 2-year follow-up. We found that patients undergoing medial UKA had a significantly higher FJS compared to patients undergoing TKA at a mean of 1.5-year follow-up. This significant difference remained at 2.4 years as well in favour of patients who had undergone medial UKA. Furthermore, our data suggest that no improvement in functional outcome is observed after 1-year follow-up since we did not note any significant changes when comparing our 1- and 2-year data of both cohorts. This last finding corresponds to the work of Pynsent [22] and Fitzgerald [7] who also reported no significant changes in PRO scores beyond 1-year follow-up of patients who underwent arthroplasty. However, Giesinger et al. [8] and Ko et al. [11] noted significant improvement when comparing their 2-year follow-up data with respectively 1-year and 6-month follow-up data. Our data might be explained by our average follow-up of 1.5 and 2.4 years following surgery and not 12 and 24 months. Lastly, we did not note a significant influence of gender on joint

 Table 2
 Forgotten Joint Score

FJS	1 year	2 year	p value
UKA	73.9	74.3	n.s.
TKA	59.3	59.8	n.s.
p value	0.002	0.004	

Note that no significant improvement was observed after 1 year of surgery for both groups

Table 3 Forgotten Joint Score by gender

	1 year	2 year
UKA		
Male	74.9	76.1
Female	73.0	72.1
p value	n.s.	n.s.
TKA		
Male	57.7	58.3
Female	59.1	60.8
<i>p</i> value	n.s.	n.s.

awareness of patients who had undergone medial UKA or TKA (Table 3).

There is a paucity of outcome studies utilizing the FJS in the literature. To our knowledge there has been only one comparative study between the UKA and TKA using the FJS. Thienpont and associates [25] found no significant differences in the FJS between patients after UKA and patients after TKA at an average of 2 years following surgery (range 1–3 years). However, in our prospective cohort of patients significantly higher scores are noted for patients who have undergone medial UKA (FJS 1 year 73.9 \pm 22.8, FJS 2 year 74.3 \pm 24.8) than patients who have undergone TKA (FJS 1 year 59.3 \pm 29.5, FJS 2 year 59.8 \pm 31.5). Since both differences are significant at both moments of follow-up, our data indicate that this difference is present at 1 year following implantation and does not improve in the time that will follow.

Comparing the baseline characteristics of both groups, we noted significant higher BMI in the TKA group $(28.6 \pm 3.7 \text{ vs. } 30.3 \pm 4.7, p = 0.02)$. The potential influence of BMI on the functional outcome, survivorship and complication rate following arthroplasty has been extensively studied. Most of the published studies have shown that obesity leads to a higher rate of infections [10, 17] and inferior implant survival [4, 6, 9]. No consensus exists, however, on the influence or relation between obesity and functional outcomes following medial UKA [16, 26] or TKA [1]. In a recent systematic review of 9 studies (670 patients), the influence of obesity on the outcome following TKA was evaluated [10]. The Knee Society Score between obese patients, defined as a BMI > 30 kg/m², and nonobese patients, defined as a BMI \leq 30 kg/m², differed significantly by 3.23 points in favour of the non-obese group. However, this minimal difference is unlikely to be clinically relevant. Baker et al. [1] recently reported the association of BMI and outcome following TKA using data from the National Joint Registry of England and Wales (pre- and post-operative questionnaires; Oxford Knee Score, EQ-5D index and EQ-5D VAS). Patients were divided by BMI in the following groups: non-obese (BMI < 25 kg/m²), obese (BMI 25–39.9 kg/m²) and morbidly obese (BMI 40–60 kg/ m^2). A total of 13,673 patients were included with an average BMI of 31.0 ± 5.5 . They found that the improvements in outcomes between the non-obese and obese patients did not differ significantly. Furthermore, the morbidly obese group had significantly lower post-operative scores than the non-obese and obese groups. Murray et al. [16] evaluated the effect of BMI on the clinical outcome of 2438 medial Oxford UKAs and found no relation between weight and outcome following medial UKA. In terms of BMI of our two cohorts, we noted a significant pre-operative difference. However, since we excluded patients with a BMI > 40 kg/ m^2 , we believe that the difference of BMI (i.e. 1.7 kg/m²) between our two cohorts is too marginal to influence outcome. Furthermore, when we stratified our data for BMI, gender, age and follow-up, the significant FJS differences remained.

With respect to registry data, the Norwegian Arthroplasty Register compared the PRO scores between TKA and UKA [13]. At a minimum follow-up of 2 years (mean, 6.5 years), 972 TKAs and 372 UKAs were compared. The outcome questionnaires being used were the Knee Injury and Osteoarthritis Outcome Score (KOOS), the EQ-5D and the visual analogue scale (VAS). The authors found that there were some significant differences in favour of the UKA group. However, the differences were too small to be considered clinically relevant. The National Registry of England and Wales compared the Oxford Knee Score (OKS) and the EQ-5D between 23,393 patients who underwent TKA and 505 patients after UKA (median followup of 6.6 months) [2]. No differences in both scores were reported. The authors highlighted that these PRO scores are unable to detect top-end differences and therefore are suboptimal measures in assessing outcomes following knee arthroplasty. Due to the ceiling effects of the traditional scores, we decided to conduct this study using the recently introduced FJS without this limitation.

Despite these results, there are several limitations to the present study. First, all procedures were performed by the senior author who has extensive experience in robotassisted UKA implantation and TKA using patient-specific guides. Therefore, results may be influenced by the subtleties specific to the respective surgical techniques and may not be duplicable in low-volume centres. Second, the use of the FJS has the advantage of not being influenced by ceiling effects. However, the score can only be used following surgery since it measures the ability of patients to forget their artificial joint in daily life. Since patients undergoing medial UKA were only affected by medial compartment OA and patients undergoing TKA by multicompartmental OA, it might be possible that they both show the same improvement following knee replacement since pre-operative differences might be present. Unfortunately, this question cannot be answered with use of the FJS. Therefore, future outcome scores, which are not limited by ceiling effects and are usable pre-operative, are needed to address these questions.

Conclusion

Our data suggest that patients who undergo UKA are better able to "forget" their artificial joint in daily life compared to patients undergoing TKA. We speculate that this observed difference may be due to the fact that UKA is a more soft-tissue and bone-conserving surgical procedure than TKA. In order to optimize the outcome of patients undergoing knee arthroplasty, this study suggests that—if possible—joint-conserving surgical strategies should be pursued.

Appendix: FJS-12 score

The following 12 questions refer to how aware you are of your artificial hip/knee joint in everyday life. Please tick one answer from each question.

Are you aware of your artificial joint...

- 1. ... in bed at night?
- never almost never seldom sometimes mostly
 2. ... when you are sitting on a chair for more than 1 h?
 never almost never seldom sometimes mostly
- 3. ... when you are walking for more than 15 min?
- never almost never seldom sometimes mostly4. ... when you are taking a bath/shower?
- never almost never seldom sometimes mostly
 5. ... when you are traveling in a car?
- O never O almost never O seldom O sometimes O mostly6. ... when you are climbing stairs?
- never almost never seldom sometimes mostly7. ... when you are walking on uneven ground?
- never almost never seldom sometimes mostly
- 8. ... when you are standing up from a low-sitting position?
- never almost never seldom sometimes mostly

9. ... when you are standing for long periods of time?
O never O almost never O seldom O sometimes O mostly
10. ... when you are doing housework or gardening?
O never O almost never O seldom O sometimes O mostly
11. ... when you are taking a walk/hiking?
O never O almost never O seldom O sometimes O mostly
12. ... when you are doing your favorite sport?
O never O almost never O seldom O sometimes O mostly

Scoring: For scoring the FJS-12, all responses are summed (never, 0 points; almost never, 1 point; seldom, 2 points; sometimes, 3 points; mostly, 4 points) and then divided into the number of completed items. This mean value is subsequently multiplied by 25 to obtain a total score range of 0–100. Finally, the score is subtracted from 100, to change the direction of the final score in a way that high scores indicate a high degree of "forgetting" the artificial joint, that is, a low degree of awareness.

If more than four responses are missing, the total score should not be used.

References

- Baker P, Petheram T, Jameson S, Reed M, Gregg P, Deehan D (2012) The association between body mass index and the outcomes of total knee arthroplasty. J Bone Joint Surg Am 94(16):1501–1508
- Baker PN, Petheram T, Jameson SS, Avery PJ, Reed MR, Gregg PJ, Deehan DJ (2012) Comparison of patient-reported outcome measures following total and unicondylar knee replacement. J Bone Joint Surg Br 94(7):919–927
- Behrend H, Giesinger K, Giesinger JM, Kuster MS (2012) The "forgotten joint" as the ultimate goal in joint arthroplasty: validation of a new patient-reported outcome measure. J Arthroplasty 27(3):430–436
- Berend KR, Lombardi AV Jr, Mallory TH, Adams JB, Groseth KL (2005) Early failure of minimally invasive unicompartmental knee arthroplasty is associated with obesity. Clin Orthop Relat Res 440:60–66
- Berger RA, Meneghini RM, Jacobs JJ, Sheinkop MB, Della Valle CJ, Rosenberg AG, Galante JO (2005) Results of unicompartmental knee arthroplasty at a minimum of 10 years of follow-up. J Bone Joint Surg Am 87(5):999–1006
- Bonutti PM, Goddard MS, Zywiel MG, Khanuja HS, Johnson AJ, Mont MA (2011) Outcomes of unicompartmental knee arthroplasty stratified by body mass index. J Arthroplasty 26(8):1149–1153
- Fitzgerald JD, Orav EJ, Lee TH, Marcantonio ER, Poss R, Goldman L, Mangione CM (2004) Patient quality of life during the 12 months following joint replacement surgery. Arthritis Rheum 51(1):100–109
- Giesinger K, Hamilton DF, Jost B, Holzner B, Giesinger JM (2014) Comparative responsiveness of outcome measures for total knee arthroplasty. Osteoarthr Cartil 22(2):184–189
- Heck DA, Marmor L, Gibson A, Rougraff BT (1993) Unicompartmental knee arthroplasty. A multicenter investigation with long-term follow-up evaluation. Clin Orthop Relat Res 286:154–159

- Kerkhoffs GM, Servien E, Dunn W, Dahm D, Bramer JA, Haverkamp D (2012) The influence of obesity on the complication rate and outcome of total knee arthroplasty: a metaanalysis and systematic literature review. J Bone Joint Surg Am 94(20):1839–1844
- 11. Ko Y, Lo NN, Yeo SJ, Yang KY, Yeo W, Chong HC, Thumboo J (2013) Comparison of the responsiveness of the SF-36, the Oxford knee score, and the knee society clinical rating system in patients undergoing total knee replacement. Qual Life Res 22(9):2455–2459
- 12. Koskinen E, Eskelinen A, Paavolainen P, Pulkkinen P, Remes V (2008) Comparison of survival and cost-effectiveness between unicondylar arthroplasty and total knee arthroplasty in patients with primary osteoarthritis: a follow-up study of 50,493 knee replacements from the finnish arthroplasty register. Acta Orthop 79(4):499–507
- Lygre SH, Espehaug B, Havelin LI, Furnes O, Vollset SE (2010) Pain and function in patients after primary unicompartmental and total knee arthroplasty. J Bone Joint Surg Am 92(18):2890–2897
- Lyons MC, MacDonald SJ, Somerville LE, Naudie DD, McCalden RW (2012) Unicompartmental versus total knee arthroplasty database analysis: is there a winner? Clin Orthop Relat Res 470(1):84–90
- Marx RG, Jones EC, Atwan NC, Closkey RF, Salvati EA, Sculco TP (2005) Measuring improvement following total hip and knee arthroplasty using patient-based measures of outcome. J Bone Joint Surg Am 87(9):1999–2005
- Murray DW, Pandit H, Weston-Simons JS, Jenkins C, Gill HS, Lombardi AV, Dodd CA, Berend KR (2013) Does body mass index affect the outcome of unicompartmental knee replacement? Knee 20(6):461–465
- Namba RS, Paxton L, Fithian DC, Stone ML (2005) Obesity and perioperative morbidity in total hip and total knee arthroplasty patients. J Arthroplasty 20(7):46–50
- Niinimaki TT, Eskelinen A, Ohtonen P, Junnila M, Leppilahti J (2012) Incidence of osteotomies around the knee for the treatment of knee osteoarthritis: a 22-year population-based study. Int Orthop 36(7):1399–1402

- Noticewala MS, Geller JA, Lee JH, Macaulay W (2012) Unicompartmental knee arthroplasty relieves pain and improves function more than total knee arthroplasty. J Arthroplasty 27(8):99–105
- Nwachukwu BU, McCormick FM, Schairer WW, Frank RM, Provencher MT, Roche MW (2014) Unicompartmental knee arthroplasty versus high tibial osteotomy: United States practice patterns for the surgical treatment of unicompartmental arthritis. J Arthroplasty 29(8):1586–1589
- Pearle AD, O'Loughlin PF, Kendoff DO (2010) Robotassisted unicompartmental knee arthroplasty. J Arthroplasty 25(2):230–237
- 22. Pynsent PB, Adams DJ, Disney SP (2005) The Oxford hip and knee outcome questionnaires for arthroplasty. J Bone Joint Surg Br 87(2):241–248
- Roche M, O'Loughlin PF, Kendoff D, Musahl V, Pearle AD (2009) Robotic arm-assisted unicompartmental knee arthroplasty: preoperative planning and surgical technique. Am J Orthop (Belle Mead, NJ) 38(2):10–15
- 24. Sun PF, Jia YH (2012) Mobile bearing UKA compared to fixed bearing TKA: a randomized prospective study. Knee 19(2):103–106
- 25. Thienpont E, Opsomer G, Koninckx A, Houssiau F (2014) Joint awareness in different types of knee arthroplasty evaluated with the Forgotten Joint score. J Arthroplasty 29(1):48–51
- Thompson SA, Liabaud B, Nellans KW, Geller JA (2013) Factors associated with poor outcomes following unicompartmental knee arthroplasty: redefining the "classic" indications for surgery. J Arthroplasty 28(9):1561–1564
- Thompson SM, Salmon LJ, Webb JM, Pinczewski LA, Roe JP (2015) Construct validity and test re-test reliability of the Forgotten Joint Score. J Arthroplasty. doi:10.1016/j.arth.2015.05.001
- Weinstein AM, Rome BN, Reichmann WM, Collins JE, Burbine SA, Thornhill TS, Wright J, Katz JN, Losina E (2013) Estimating the burden of total knee replacement in the United States. J Bone Joint Surg Am 95(5):385–392